

POULTRY FEEDS

1310

PRODUCT DATA

PRODUCT DATA		
ME	. 1443.46	Kcal/lb
PROTEIN	. 11.139	%
ARGININE	. 0.397	%
CYSTINE	. 0.111	%
HISTIDINE	0.203	%
ISOLEUCINE		%
LEUCINE		%
LYSINE	. 0.495	%
METHIONINE		%
PHENYLALANINE	. 0.433	%
THREONINE	. 0.314	%
TRYPTOPHAN	. 0.088	%
TYROSINE		%
VALINE	. 0.411	%
FAT	5.090	%
LINOLEIC ACID		%
FIBER	. 3.975	%
FIBERCALCIUM	. 0.710	%
PHOSPHORUS	. 0.415	%
AVAIL. PHOSPH	. 0.062	%
COPPER		mg/lb
IODINE	. 0.017	mg/lb
IRON		mg/lb
MAGNESIUM		%
MANGANESE	. 7.102	mg/lb
POTASSIUM	. 0.546	%
SELENIUM		mg/lb
ZINC	9.072	mg/lb
BIOTIN	0.066	mg/lb
CHOLINE		mg/lb
FOLIC ACID		mg/lb
NIACIN	. 14.638	mg/lb
PANTOTHENIC ACID	4.265	mg/lb
PYRIDOXINE	. 2.773	mg/lb
RIBOFLAVIN	0.643	mg/lb
THIAMINE		mg/lb
VITAMIN E	7.876	IU/lb

TFP NUTRITION NACOGDOCHES, TX

SELECT SCRATCH


A mixture of select high quality grains, protein, grit and oils to supplement the daily diet of chickens.

FEATURES

- * High Energy 1443 Kcal/lb
- * Corn, grain sorghum, & oats
- * Black Oil Sunflower Seeds
- * Protein pellets
- * Oyster shell grit
- * Vegetable oil

BENEFITS

- * Feed as a scratch grain to encourage natural pecking and foraging instincts.
- * Helps maintain good flock condition
- Oyster shell provides grit needed for feed digestion
- Sunflower seeds and vegetable oil provide excellent sources of fat



GUARANTEED ANALYSIS

Crude Protein, minimum	0.40% 0.20% 5.00% 5.00% 0.50% 1.00% 0.30%
Salt (NaCl), minimum Salt (NaCl), maximum	0.10%

INGREDIENTS

Corn chops, grain sorghum, oats, black oil sunflower seeds, oyster shell, cane molasses, vegetable oil, soybean meal, wheat middlings, calcium carbonate, monocalcium phosphate, salt, natural and artificial flavors.

FEEDING DIRECTIONS

Feed as a scratch grain supplement to adult chickens at a rate not to exceed 10% of the diet. This product is not a complete feed and should be fed along with Lone Star 20% Protein Laying Mash Pellets. Provide a source of fresh, clean water at all times.

CAUTION: Feed should be stored in a well-ventilated, dry area protected from rodents and insects. Feed is perishable. Do not feed moldy or insect infested feed to birds as it may cause illness or death.